

Special Protocol

(Effective through period of Commissioner's Order)

Administration of Immunizations by EMT-Basics Pursuant to a Public Health Order of the Commissioner: COVID-19 VACCINE

Vaccines for the novel Coronavirus 2019 (COVID-19), caused by the virus SARS-CoV-2, are given by injection into the muscle (IM). EMT-Basics are not trained to administer intramuscular medications, nor previously ever authorized to administer vaccinations. EMT-Basics operate in accordance with the Department's EMS System regulations and Statewide Treatment Protocols of the MDPH Office of Emergency Medical Services (OEMS), and in connection with an MDPH-licensed ambulance service. They also obtain authorization to practice from their ambulance service's affiliate hospital medical director.

The specific authorization granted by the latest Commissioner's Order, pursuant to the Department's DCP regulations, 105 CMR 700.004(H), authorizes currently certified Massachusetts EMT-Basics working in connection with an ambulance service to administer vaccine for the prevention of COVID-19, in persons covered by a vaccine emergency use authorization (EUA), in accordance with this Special Protocol issued by OEMS. Prior to being deployed by their ambulance service to administer COVID-19 vaccine, EMT-Basics must have successfully completed an IM vaccination training program as specified below.

Minimum Requirements: IM vaccination training for EMT-Basics

Any EMT-Basic who is administering IM COVID-19 vaccination in conjunction with their ambulance service must receive training from their Affiliate Hospital Medical Director or designee.

Such training shall include:

1. Sterile technique
2. Familiarization with the equipment needed
3. How to choose needle and syringe
4. Familiarization with required consent and record keeping
5. Indications and contraindications to the vaccine in question
6. How to draw up vaccine safely and sterilely
7. How to select and prep injection site
8. How to give the injection – follow attached CDC documents, listed under Reference, below
9. Post injection site care
10. Post injection patient observation
11. Familiarization with instructions to give patient after vaccination

Total expected time of instruction: 4 hours

Ambulance service must maintain records of such training.

Reference: <https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/administration.html>

Two PDF documents are available on the OEMS website as references as well: "CDC IM-injection-adult" and "CDC ACIP Vaccine Administration".

Deployment

All EMT-Basics vaccinating under this waiver must be working with at least one physician, nurse, or paramedic **immediately available** at the treatment site as the supervisor of dosing, procedure, and technique.

EMT-Basics may not thaw or reconstitute vaccine. They are limited to drawing up doses and administering them to recipients.

EMT-Basics must be trained and authorized by their AHMD or designee prior to administering COVID-19 vaccine in any form. Under this Special Protocol, EMT-Basics can only administer vaccine while working for their ambulance service.

Such training must also include the procedures to be followed by EMT-Basics within the setting in which the vaccines are to be administered, for determining type and dose of vaccine to be given, including the identity of staff who will make such determination.

EMT-BASIC STANDING ORDERS

- a. Provide patient, parent or legal representative with a copy of the Vaccine EUA Fact Sheet and answer any questions.
- b. Screen for contraindications according to accompanying Special Protocol Appendix A1, and/or the CDC/ACIP document (or ascertain the result of such screening)
- c. Always check the package insert or procedural instructions prior to administration of any vaccine. Follow insert or procedural instructions to prepare and draw up vaccine dose.
- d. Administer IM vaccines at a 90° angle with 22-25-gauge needle. The needle length for IM injections depends upon the age, gender, and/or weight of the vaccine recipient. (See Table attachment.). Administer IM, according to the recommended age-specific dose and schedule. (See Special Protocol Appendix A1.)
- e. Administer vaccine.
- f. Observe patient for an allergic reaction for 15-20 minutes after administering vaccine. If an anaphylactic/allergic reaction occurs, treat according to Protocol 2.2A/2.2P Allergic Reaction/Anaphylaxis.
- g. Report clinically significant adverse events to the Vaccine Adverse Event Reporting System (VAERS) at 1-800-822-7967 or <http://www.vaers.hhs.gov/>.

Special Protocol

(Effective through period of Commissioner's Order)



STANDING ORDERS

Needle Length and Injection Site for IM Injection

Gender and Weight	Needle Length	Injection Site	Injection Technique
Children (16-18 years)	5/8" - 1"	Deltoid	Depends on Body Mass
Male and Female <60 kg (<130 lbs)	5/8"	Deltoid	Do not bunch subcutaneous and muscle tissue
Male and Female (130-152 lbs)	1"		
Female 70-90kg (152-200 lbs)	1"-1 1/2"		
Male 70kg-118kg (152-260 lbs)			
Female >90kg (200lbs)	1 1/2"		
Male >118kg (260 lbs)			

E